


## Recombinant human MMP13 protein

Catalog#:P05786    Derived from E.coli

<b>DESCRIPTION</b>	Recombinant Human MMP13 protein is produced by our E.coli expression system with His tag. Uniprot: P45452 Gene ID: 4322
<b>Size</b>	40 kDa
<b>FORMULATION</b>	Lyophilized from a 0.2 µm filtered solution of Tris-HCl, PH 8.0.
<b>SHIPPING</b>	The product is shipped at -20°C temperature. Upon receipt, store it immediately at the temperature listed below.
<b>STORAGE</b>	Reconstituted protein solution can be stored at 4-7°C for 2-3 months, stable at < -20°C for 1-2 years.
<b>RECONSTITUTION</b>	Reconstituted protein solution can be diluted with distilled PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>QUALITY CONTROL</b>	(Liquid)Concentration: 1 mg/mL as determined by BCA. Purity: 85% as determined by reducing SDS-PAGE.
<b>AMINOACID</b>	Recombinant protein is produced by our E.coli expression system and the target gene encoding 1-350aa is expressed with a 6*His tag at the N-terminus.
<b>BACKGROUND</b>	Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CCN2. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CCN2. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.



Recombinant human MMP13 protein