

Recombinant Human NOL3

Catalog#:P01232 Derived from *E.coli*

DESCRIPTION	<p>Recombinant Human Nucleolar Protein 3 is produced by our E.coli expression system and the target gene encoding Met1-Ser208 is expressed.</p> <p>Accession#: O60936</p> <p>Known as: Nucleolar Protein 3; Apoptosis Repressor With CARD; Muscle-Enriched Cytoplasmic Protein; Myp; Nucleolar Protein of 30 kDa; Nop30</p>
FORMULATION	<p>Supplied as a 0.2 μm filtered solution of 25mM Tris-HCl, 1mM DTT, 1mM EDTA, 2mM β-ME, 20% Glycerol, pH 7.5.</p>
SHIPPING	<p>The product is shipped on dry ice/polar packs.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Store at ≤-70°C, stable for 6 months after receipt.</p> <p>Store at ≤-70°C, stable for 3 months under sterile conditions after opening.</p> <p>Please minimize freeze-thaw cycles.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass:22.63kDa AP Mol Mass:29kDa, reducing conditions.</p> <p>Purity: Greater than 90% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>Nucleolar protein 3 is encoded by NOL3 gene. Multiple transcript variants encoding different isoforms have been found for this gene. So far, Nucleolar protein 3 has show to have two Isoforms. Isoform 1 may be involved in RNA splicing. Isoform 2 functions as an apoptosis repressor that blocks multiple modes of cell death. It inhibits extrinsic apoptotic pathways through two different ways. Firstly, it by interacting with FAS and FADD upon FAS activation blocking death-inducing signaling complex (DISC) assembly. Secondly by interacting with CASP8 in a mitochondria localization- and phosphorylation-dependent manner, limiting the amount of soluble CASP8 available for DISC-mediated activation. It has been shown to down-regulate the enzyme activities of caspase 2, caspase 8 and tumor protein p53.</p>