

Recombinant Human CEACAM5

Catalog#:P01058 Derived from Human Cells

DESCRIPTION	<p>Recombinant Human Carcinoembryonic Antigen- Related Cell Adhesion Molecule 5 is produced by our Mammalian expression system and the target gene encoding Lys35-Ala685 is expressed with a 6His tag at the C-terminus.</p> <p>Accession#: NP_004354.3</p> <p>Known as: Carcinoembryonic antigen- related cell adhesion molecule 5; Carcinoembryonic antigen; CEA; Meconium antigen 100; CD66e; CEACAM5</p>
FORMULATION	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at $<-20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at $<-20^{\circ}\text{C}$ for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100$\mu\text{g/ml}$.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass: 72.4kDa AP Mol Mass: 95-150kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>Carcinoembryonic antigen-related cell adhesion molecules (CEACAMs) belong to a group of mammalian immunoglobulin related glycoproteins. They play critical roles in cell-cell recognition. CEACAM5, also called CEA and CD66e, is characterized by having seven extracellular Ig domains and a glycosylphosphatidylinositol (GPI) anchor. CEACAM5 is expressed primarily by epithelial cells, and functions as a calcium-independent adhesion molecule through homophilic and heterophilic interactions with CEACAM1. Studies have shown that CEACAM5 is overexpressed in a majority of carcinomas, and its overexpression can protect tumor cells from apoptosis. It is commonly used as a cancer marker.</p>

