

Tel: 400-968-6088 Fax: 010-56371281

Http://www.solarbio.com

## **Recombinant Mouse TSLP**

Catalog#:P00835 Derived from Human Cells

Catalog#.1 00055 Derived from Human Cens	
DESCRIPTION	Recombinant Mouse Thymic stromal lymphopoietin is produced by our
	Mammalian expression system and the target gene encoding Tyr20-Glu140 is expressed with a Fc tag at the C-terminus.
	Accession#: Q9JIE6
	<b>Known as</b> : Thymic stromal lymphopoietin; Thymic stroma-derived lymphopoietin; Tslp
	<del>'                                   </del>
FORMULATION	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
SHIPPING	The product is shipped at ambient temperature.
	Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at <-20°C, though stable at room
	temperature for 3 weeks.
	Reconstituted protein solution can be stored at 4-7°C for 2-7 days.
	Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.
	It is not recommended to reconstitute to a concentration less than 100µg/ml.
	Dissolve the lyophilized protein in distilled water.
	Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY	Mol Mass:41.1kDa AP Mol Mass:42-58kDa, reducing conditions.
	<b>Purity</b> : Greater than 95% as determined by reducing SDS-PAGE.
CONTROL	Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	Thymic stromal lymphopoietin (TSLP) is a protein belonging to the cytokine family, contains 140 amino acids. It is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. TSLP induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c+ dendritic cells. It can induce
	particular, enhances the maturation of CD11c+ dendritic cells. It can induce allergic inflammation by directly activating mast cells. TSLP is produced mainly by non-hematopoietic cells such as fibroblasts, epithelial cells and different types of stromal or stromal-like cells. These cells are located in regions where TSLP activity is required.
Q2, 350 - 1907 A 20 - 1907 A 2	

