

Tel: 400-968-6088 Fax: 010-56371281

Http://www.solarbio.com

Recombinant Human CD79B

Catalog#:P00542 Derived from Human Cells

Editiogh: 100312 Bettved from Human Cens	
DESCRIPTION	Recombinant Human CD79B is produced by our Mammalian expression system and the target gene encoding Ala29-Asp159 is expressed with a Fc tag at the
	C-terminus.
	Accession#: P40259
	Known as: B-Cell Antigen Receptor Complex-Associated Protein Beta Chain;
	B-Cell-Specific Glycoprotein B29; Ig- Beta; Immunoglobulin-Associated B29
	Protein; CD79b; CD79B; B29; IGB
FORMULATION	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
SHIPPING	The product is shipped at ambient temperature.
	Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at<-20°C, though stable at room
	temperature for 3 weeks.
	Reconstituted protein solution can be stored at 4-7°C for 2-7 days.
	Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.
	It is not recommended to reconstitute to a concentration less than 100µg/ml.
	Dissolve the lyophilized protein in distilled water.
	Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY	Mol Mass:42.1kDa AP Mol Mass:45-65kDa, reducing conditions.
	Purity : Greater than 95% as determined by reducing SDS-PAGE.
CONTROL	Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	CD79B is a single-pass type I membrane protein. CD79B contains one Ig-like V-type domain and one ITAM domain. CD79B is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR), which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79B enhances phosphorylation of CD79A, possibly by recruiting kinases that phosphorylate CD79A or by recruiting proteins that bind to CD79A and protect it from dephosphorylation.
	CD79A for initiation of the signal transduction cascade activated by the B-cell
	antigen receptor complex (BCR), which leads to internalization of the complex,
	trafficking to late endosomes and antigen presentation. CD/9B enhances
	CD79A or by recruiting proteins that bind to CD79A and protect it from
	dephosphorylation.
10. 10.	



