

Recombinant Human IGFBP-5

Catalog#:P02272 Derived from Human Cells

DESCRIPTION	<p>Recombinant Human Insulin- Like Growth Factor-Binding Protein 5 is produced by our Mammalian expression system and the target gene encoding Leu21-Glu272 is expressed with a 6His tag at the C-terminus.</p> <p>Accession#: P24593</p> <p>Known as: Insulin- Like Growth Factor-Binding Protein 5; IBP-5; IGF- Binding Protein 5; IGFBP-5; IGFBP5;IBP5</p>
FORMULATION	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at <-20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100μg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass:29.61kDa AP Mol Mass:30-40kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>Insulin- Like Growth Factor-Binding Protein 5 (IGFBP-5) is a secreted protein that belongs to the insulin-like growth factor (IGF) binding proteins superfamily. Members of this family prolong the half-life of the IGFs. They have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors. IGFBP-5 contains one IGFBP N- terminal domain and one thyroglobulin type- 1 domain. IGFBP-5 is expressed by fibroblasts, myoblasts and Osteosarcoma. It is also present at lower levels in liver, kidney, and brain.</p>
SDS-PAGE	