

Tel: 400-968-6088 Fax: 010-56371281

Http://www.solarbio.com

Recombinant Human CXCL10

Catalog#:P01205 Derived from Human Cells

DESCRIPTION	Recombinant Human C-X-C Motif Chemokine 10 is produced by our Mammalian expression system and the target gene encoding Val22-Pro98 is expressed with a human IgG1 Fc, 6His tag at the C-terminus. Accession#: P02778 Known as: C-X-C Motif Chemokine 10; 10 kDa Interferon Gamma-Induced Protein; Gamma-IP10; IP-10; Small-Inducible Cytokine B10; CXCL10; INP10; SCYB10
FORMULATION	Lyophilized from a 0.2µm filtered solution of PBS, pH 7.4.
SHIPPING	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY CONTROL	Mol Mass:36.6kDa AP Mol Mass:35kDa, reducing conditions. Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	Human C-X-C Motif Chemokine Ligand 10 (CXCL10) is a non-ELR chemokine secreted by various cell types, such as monocytes, endothelial cells and fibroblasts, in response to IFN-γ. CXCL10 functions via chemokine receptor CXCR3. CXCL10 has been attributed to several roles, such as chemoattraction for activated T-lymphocytes, inhibition of angiogenesis, and antitumor activity.

