

Tissue Inorganic Phosphorus Content Assay Kit

Note: Take two or three different samples for prediction before test.

Detection instrument: Spectrophotometer

Cat No: BC2840

Size: 50T/48S

Components:

Reagent I: 50 mL×1. Storage at 4°C.

Reagent II: 10 mL×1. Storage at 4°C.

Reagent III: Powder×2. Storage at 4°C. Fully dissolve with 10mL of distilled water, then add 5 mL of Reagent II and mix well.

Standard: 1 mL×1, 1 mmol/L inorganic phosphorus standard. Storage at 4°C.

Product Description:

Inorganic phosphorus mainly refers to phosphate radical, which is involved in many kinds of metabolism, including energy metabolism, nucleic acid metabolism, protein phosphorylation and dephosphorylation, and so on. In addition, inorganic phosphorus promotes the synthesis, transformation and transport of carbohydrates.

Molybdenum blue can react with phosphate group, the reaction product can be detected by colorimetric assay at 660nm and calculate the inorganic phosphorus content indirectly.

Required material

Spectrophotometer, centrifuge, water bath, 1mL glass cuvette, transferpette, homogenizer/mortar, distilled water.

Procedure:

I. Sample Extraction:

Suggest 0.1g of sample with 1mL of Reagent I, fully grind on ice, centrifuge at 10000 rpm and 4°C for 10 minutes, supernatant is used for test.

II. Determination Procedure:

1. Preheat the spectrophotometer for 30min, adjust wavelength to 660nm, set zero with distilled water.
2. Set the temperature of water bath to 40°C.
3. Add reagents with the following list:

Reagent name (μL)	Blank tube (A _B)	Standard tube (A _T)	Test tube (A _S)
Standard		50	
Supernatant			50
Distilled water	500	450	450
Reagent III	500	500	500

Mix well, 40°C water bath for 10 minutes, detect the absorbance at 660 nm after cooling at room temperature for 10 minutes. Record as A_B , A_S and A_T respectively.

Note: The colorimetry needs to be completed within 40 minutes.

Calculation:

$$\text{Phosphate (mmol/g fresh weight)} = [C \times (A_T - A_B) \div (A_S - A_B)] \times V \div W = 0.001 \times (A_T - A_B) \div (A_S - A_B) \div W$$

C: standard concentration, 1 mmol/L;

V: supernatant volume, 1 mL=0.001 L;

W: Sample weight, g

Note:

1. Reagent III should be prepared before use and only be used the same day.
2. Before the determination, 1-2 samples shall be used for pre-test. If the absorption value is greater than 0.8, distilled water shall be used for corresponding dilution.

Experimental example:

1. 0.1g liver is added with 1 mL of Reagent I, and the supernatant is centrifuged. Then, the determination procedure is followed. $A_T = 0.789$, $A_B = 0.016$, $A_S = 0.429$.

$$\text{Inorganic phosphorus content (mmol/g mass)} = 0.001 \times (A_T - A_B) \div (A_S - A_B) \div W = 0.001 \times (0.789 - 0.016) \div (0.429 - 0.016) \div 0.1 = 0.019 \text{ mmol/g mass.}$$

Related Products:

Bu J, Yu J, Wu Y, et al. Hyperlipidemia affects tight junctions and pump function in the corneal endothelium[J]. The American Journal of Pathology, 2020.

Related Products:

BC2860/BC2865 Serum Total Iron Binding Capacity(TIBC) Assay Kit

BC2850/BC2855 Total Phosphorus Content Assay Kit

BC4350/BC4355 Tissue Iron Content Assay Kit

BC4380/BC4385 Blood Ammonia Content Assay Kit

Technical Specifications:

The detection limit: 0.0576 mmol/L

Linear range: 0.0625-2 mmol/L