

苏丹III染色液

货号: G1510 **规格:** 100mL

保存: 室温, 避光保存, 有效期1年。

产品介绍:

中性脂肪(Neutral fat)是由三分子脂肪酸和一分子甘油组成的脂类,呈中性。中性脂肪是储存能量的方式之一,在氧化时释放出能量。在正常情况下,除脂肪细胞外,其他细胞在光学显微镜下几乎看不到脂滴,如果细胞质内出现大量脂滴即为脂肪变性,常见于肝细胞、心肌细胞、肾曲管上皮细胞等。中性脂肪染色经常采用苏丹II、苏丹III、苏丹IV、苏丹黑 B、油红 O 法等。苏丹染料脂质染色的机理一般认为纯属物理学的脂溶作用和吸附作用。苏丹类染料由于在脂质中的溶解度大于在有机溶剂的溶解度,所以染色时染料便从染液中转移到被染的脂质中去,使脂质呈现出染液的颜色。

苏丹Ⅲ染色液主要用于显示组织器官的脂肪变性和类脂质的异常沉着,常发生于肝、肾、心等实质脏器的脂肪变性,细胞内出现多数中性脂肪滴;鉴别和诊断脂肪组织中所发生的肿瘤及其性质。标本不采用含有乙醇的固定液(如需固定可采用 10%中性福尔马林)、也不采用石蜡切片,需用冰冻切片或碳蜡切片。

自备材料:

载玻片、显微镜、70%乙醇、蒸馏水、S2150-甘油明胶封片剂或 S2155-水性封片剂(PVP)

操作步骤: (仅供参考)

- 1. 新鲜组织低温切片,一般-20℃到-25℃。如样本为脂肪瘤,应调节至-30℃。
- 2. 冰冻切片 6-15μm(6-8μm 为佳), 贴于载玻片上。70%乙醇稍微浸洗一下。
- 3. 入苏丹III染色液,浸染 10min。70%乙醇洗去多余染液。蒸馏水浸洗 1min。
- 4. 按实验具体要求进行下游实验。
- 5. 甘油明胶或 S2155-水性封片剂 (PVP) 封固。


染色结果:

# 1 d	中性脂肪	橘红色	N.

注意事项:

- 1. 标本不宜采用含有乙醇的固定液、也不宜用石蜡切片,需用冰冻切片。如需固定应采用 10%中性福尔 马林固定液或 10%甲醛-钙液。
- 2. 本试剂为过饱和溶液,建议临用前过滤使用。
- 3. 在染色过程中必须防止染料发生沉淀。故切片入染液时应密封,勿与流动空气相接触,避免溶液挥发时发生沉淀。
- 4. 冰冻切片较易着色, Mayer 苏木素复染时应避免过染。
- 5. 苏丹染料容易褪色,应密闭保存。
- 6. 甘油明胶封固的样本,保存时间不长。如需长期保存,可以在盖玻片与载玻片交界的边缘用中性树胶 封闭。
- 7. 为了您的安全和健康,请穿实验服并戴一次性手套操作。



















Sudan III Stain Solution

Cat: G1510 **Size:** 100mL

Storage: RT, avoid light, valid for 1 year.

Introduction

Neutral fat is a lipid composed of three molecules of fatty acids and one molecule of glycerol, which is neutral. Neutral fats are one of the ways to store energy, releasing energy during oxidation. Under normal circumstances, except for adipocytes, other cells can hardly see lipid droplets under an optical microscope. If a large number of lipid droplets appear in the cytoplasm, it is considered steatosis, commonly seen in liver cells, myocardial cells, renal tubular epithelial cells, etc. Neutral fat staining often uses methods such as Sudan II, Sudan III, Sudan IV, Sudan Black B, and Oil Red O. The mechanism of Sudan dye lipid staining is generally believed to be purely physical, involving lipid solubility and adsorption. Sudan dyes, due to their higher solubility in lipids than in organic solvents, transfer the dye from the dye solution to the dyed lipids during dyeing, resulting in the lipids presenting the color of the dye solution.

Sudan III Stain Solution is mainly used to show fatty degeneration of tissues and organs and abnormal lipid-like sedation. It often occurs in fatty degeneration of liver, kidney, heart and other parenchymal organs that most of the neutral fat droplets appear in cells. It can identify and diagnose tumors and their properties in adipose tissue. The sample should not fixed with ethanol (10% formalin if required) or paraffin. Frozen sections or carbon wax sections were needed. The positive staining results of fat were orange to red, but the specific color depended on the concentration of lipid.

Self Provided Materials

Slide, Microscope, 70% alcohol, Distilled Water, S2150-Glycerol Gelatin or S2155-Aqueous Mounting Medium (PVP)

Protocol (for reference only)

- 1. Prepare low-temperature sections of fresh tissue, generally at -20°C to -25°C. If the sample is lipoma, it should be adjusted to 30 °C.
- 2. Cut the frozen section at 6-15um (6-8 um is preferable) and attach to the slide. Wash with 70% alcohol for seconds.
- 3. Soak the section in Sudan III Solution for 10min. Wash with 70% alcohol to remove excess solution. Wash with distilled water for 1 min.
- 4. Conduct downstream experiments according to specific experimental requirements.
- 5. Glycerol gelatin or S2155-Aqueous Mounting Medium(PVP) sealing.

Result

37 . 10.	O D 1
I Neutral tat	l Orange Red
1 (Catial lat	L Grange Rea

Note

- 1. Samples should not be fixed with ethanol or embedded with paraffin, and frozen sections should be used. It is recommended to use 10% neutral formalin or 10% formaldehyde-calcium solution to fix.
- 2. Precipitation of dyes must be prevented during dyeing. Therefore, slices should be covered when they are put into dye solution, so as not to contact with flowing air to avoid precipitation when the solution volatilizes.
- 3. This reagent is a supersaturated solution, it is recommended to filter it before use.
- 4. Frozen sections are easier to be colored, and overdyeing should be avoided when re-dyeing with Mayer hematoxylin.
- 5. Sudan dyes fade easily and should be kept in airtight condition.
- 6. Samples sealed with glycerol gelatin did not last long. If long-term preservation is required, the edge of the boundary between the cover slide and the slide can be sealed with neutral gum.
- 7. For your safety and health, please wear lab clothes and disposable gloves.





