

Pyruvate Carboxylase(PC) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Microplate Reader

Catalog Number: BC0735

Size: 100T/96S

Product Composition: Before use, please carefully check whether the volume of the reagent is consistent with the volume in the bottle. If you have any questions, please contact Solarbio staff in time.

Reagent Name	Size	Preservation Condition
Extract solution	Liquid 110 mL×2	2-8°C
Reagent I	Liquid 15 mL×1	2-8°C
Reagent II	Liquid 5 mL×1	2-8°C
Reagent III	Powder×1	-20°C
Reagent IV	Powder×1	-20°C
Reagent V	Liquid 2 mL×1	2-8°C
Reagent VI	Powder×1	-20°C
Reagent VI Diluent	Liquid 5 mL×1	2-8°C

Solution Preparation:

1. Reagent III: Dissolve with 3 mL of distilled water before using. It can be stored at -20°C for 2 weeks after subassembly to avoid repeated freezing and thawing.

2. Reagent IV: Dissolve with 2 mL of distilled water before using. It can be stored at -20°C for 2 weeks after subassembly to avoid repeated freezing and thawing.

3. Reagent VI: Dissolve with 0.13 mL of distilled water before using. It can be stored at -20°C for 4 weeks after subassembly to avoid repeated freezing and thawing.

4. Preparation of Reagent VI working liquid: Before use, according to the sample number, the ratio of reagent VI : Reagent VI Diluent Solution = 0.01mL : 0.3mL (total 0.31mL, about 15T) is prepared.

5. Preparation of working liquid: Before use, the working liquid was prepared according to the sample size in the ratio of reagent II: reagent III : reagent IV = 2 : 1 : 1.

Product Description:

Pyruvate carboxylase (PC, EC 6.4.1.1) is widely present in mitochondria of animals, molds and yeast, but is not found in plants and most bacteria. PC is the main postreaction for oxaloacetate, and is the first rate-limiting enzyme in the gluconeogenesis process.

PC irreversibly catalyzes pyruvate, ATP, CO₂ and water to oxaloacetate, ADP and Pi, malic dehydrogenase further catalyzes the formation of malic acid and NAD⁺ from acetoacetic acid and NADH. The enzyme activity of PC can be reflected by detecting the oxidation rate of NADH at 340 nm.

Reagents and Equipment Required but Not Provided:

Ultraviolet spectrophotometer/microplate reader, desk centrifuge, water bath, adjustable pipette, micro quartz cuvette/96 well flat-bottom UV plate, mortar/homogenizer, ice and distilled water.

Operation procedure:

I. Sample preparation(The sample size to be tested can be adjusted appropriately, and the specific proportion can be referred to the literature.)

- 1) Collecting 0.1 g of tissue or 5 million cells, add 1 mL of Extract solution, grinding on ice with mortar/homogenizer.
- 2) Centrifuge at 1000 ×g for 10 minutes at 4°C,
- 3) Take the supernatant to other tube and centrifuge at 11000 ×g for 15 minutes at 4°C
- 4) The supernatant is used to detect PC that leaking from mitochondria, which shows the effect of mitochondrial extraction.
- 5) Add 1 mL of Extract solution to the sediment, splitting with ultrasonic (power 200w, work time 5s, interval 10s, repeat 12 times), used to detect the enzyme activity of PC and protein content.

II. Determination procedure:

- 1) Preheat ultraviolet spectrophotometer/microplate reader for 30 minutes, adjust the wavelength to 340 nm, set zero with distilled water.
- 2) Preheat Reagent I at 37°C for 15 minutes.
- 3) Operation table:

Reagent (μL)	Blank tube (B)	Test tube (T)
Reagent I	90	90
Working solution	64	64
Reagent V	16	16
Reagent VI- working solution	20	20
Sample	-	10
Distilled water	10	-

Add the above reagents to the micro quartz cuvette/96 well flat-bottom UV plate in order. Mix thoroughly. Detect the absorbance at 340 nm at the time of 10 seconds record as A_{T1} or A_{B1} . Then place dishes with the reaction solution in a 37°C water bath or incubator for 2 minutes. Take it out and wipe it clean, immediately measure the absorbance at the time of 130 seconds which record as A_{T2} or A_{B2} . $\Delta A_T = A_{T1} - A_{T2}$, $\Delta A_B = A_{B1} - A_{B2}$, $\Delta A = \Delta A_T - \Delta A_B$. The blank tube only need to test once or twice.

III. PC enzyme activity calculation:

A. micro quartz cuvette

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the production of 1 nmol of NADH per minute every milligram of protein.

$$PC \text{ Activity (U/mg prot)} = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^9] \div (V_s \times C_{pr}) \div T = 1607 \times \Delta A \div C_{pr}$$

ϵ : NADH molar extinction coefficient, 6.22×10^3 L/mol/cm;

d : Light path of cuvette, 1 cm;

V_{rv} : Total reaction volume, 2×10^{-4} L;

V_s : Sample volume (mL), 0.01 mL;

C_{pr} : Sample protein concentration (mg/mL);

T : Reaction time (min), 2 minutes;

10^9 : 1 mol = 10^9 nmol.

B. 96 well flat-bottom UV plate

Change the d -1 cm in the above formula to d -0.6 cm for calculation.

Note:

1. Take one or two different samples for prediction before test. It is recommended to dilute the crude enzyme solution with the Extract solution before the determination if the $\Delta A > 0.8$. While, extending the response time (5 minutes or 10 minutes) if $\Delta A < 0.01$.
2. The blank tube is a detection hole for detecting the quality of each reagent component, and normally that the change of ΔA_B does not exceed 0.05.
3. The protein concentration of the sample needs to be determined by yourself. Since the Extract solution contains a relatively high protein concentration (about 1 mg/mL), the protein concentration of the Extract solution must be deducted when measuring the protein concentration of the sample.
4. It is recommended to use the sample protein concentration to calculate the enzyme activity. If the sample fresh weight is used to calculate, the enzyme activity of cytoplasmic extract needs to be measured, and the sum of supernatant and precipitation enzyme activity is the total enzyme activity.
5. Reagents in this kit are sufficient to complete 100 tube reactions.
6. Appendix: calculation formula of sample weight: (sample test number is 100T/48S)

A. micro quartz cuvette
1) Calculation of PC vitality in supernatant:

Unit definition: One unit of enzyme activity is the amount of enzyme catalyzes the production of 1 nmol of NADH per minute every gram of tissue.

$$\text{PC Activity (U/g weight)} = [\Delta A1 \times V_{rv} \div (\epsilon \times d) \times 10^9] \div (W \div V_e \times V_s) \div T = 1607 \times \Delta A1 \div W$$

$\Delta A1$: Supernatant absorbance;

ϵ : NADH molar extinction coefficient, 6.22×10^3 L/mol/cm;

d : Light path of cuvette, 1 cm;

V_{rv} : Total reaction volume, 2×10^{-4} L;

V_s : Sample volume (mL), 0.01 mL;

V_e : Extraction solution, 1 mL;

C_{pr} : Sample protein concentration (mg/mL);

T : Reaction time (min), 2 minutes;

10^9 : 1 mol = 10^9 nmol;

W : Sample weight, g.

2) Calculation of PC activity in precipitation:

Unit definition: One unit of enzyme activity is the amount of enzyme catalyzes the production of 1 nmol of NADH per minute every gram of tissue.

$$\text{PC Activity (U/g weight)} = [\Delta A2 \times V_{rv} \div (\epsilon \times d) \times 10^9] \div (W \div V_e \times V_s) \div T = 1607 \times \Delta A2 \div W$$

$\Delta A2$: Sediment absorbance;

ϵ : NADH molar extinction coefficient, 6.22×10^3 L/mol/cm;

d : Light path of cuvette, 1 cm;

V_{rv} : Total reaction volume, 2×10^{-4} L;

V_s : Sample volume (mL), 0.01 mL;

Ve: Sediment heavy suspension volume, 1 mL;

Cpr: Sample protein concentration (mg/mL);

T: Reaction time (min), 2 minutes;

10^9 : 1 mol=10⁹ nmol;

W: Sample weight, g.

3) Calculation of total activity of sample PC:

Total activity is the sum of PC activity in supernatant and sediment.

PC (U/g weight)= $1607 \times \Delta A_1 \div W + 1607 \times \Delta A_2 \div W$.

B. 96 well flat-bottom UV plate

Change the d-1cm in the above formula to d-0.6cm for calculation.

Experimental example:

1. 1 mL of Extract solution is added to 0.1 g of rabbit heart tissue for homogenization. The supernatant is diluted 100 times with Extract solution, and the precipitation was diluted 4 times.

Then, measured by micro quartz plate according to the determination steps, Supernatant: the $\Delta A_T = A_{1T} - A_{2T} = 1.0091 - 0.7487 = 0.2604$, $\Delta A_B = A_{1B} - A_{2B} = 0.9948 - 0.9678 = 0.027$, $\Delta A_1 = \Delta A_T - \Delta A_B = 0.2604 - 0.027 = 0.2334$, Precipitate: $\Delta A_T = A_{1T} - A_{2T} = 1.08 - 0.6157 = 0.4643$, $\Delta A_B = A_{1B} - A_{2B} = 0.9948 - 0.9678 = 0.027$, $\Delta A_2 = \Delta A_T - \Delta A_B = 0.4643 - 0.027 = 0.4373$

Supernatant: the activity of PC (U/g mass) = $1607 \times \Delta A_1 \div W \times 100$ (dilution ratio) = $1607 \times 0.2334 \div 0.1 \times 100 = 375073.8$ U/g mass;

Precipitation: the enzyme activity of PC (U/g mass) = $1607 \times \Delta A_2 \div W \times 4$ (dilution ratio) = $1607 \times 0.4373 \div 0.1 \times 4 = 28109.64$ U/g mass;

The total enzyme activity of PC (U/g mass) = $1607 \times \Delta A_1 \div W \times 100$ (dilution) + $1607 \times \Delta A_2 \div W = 1607 \times 0.2334 \div 0.1 \times 100 + 1607 \times 0.4373 \div 0.1 \times 4 = 403183.44$ U/g mass.

Recent Product Citations:

- [1] Guo X, Jiang X, Chen K, Liang Q, Zhang S, Zheng J, Ma X, Jiang H, Wu H, Tong Q. The Role of Palmitoleic Acid in Regulating Hepatic Gluconeogenesis through SIRT3 in Obese Mice. *Nutrients*. 2022 Apr 1;14(7):1482. doi: 10.3390/nu14071482. PMID: 35406095; PMCID: PMC9003329.
- [2] Chen J, Zhang J, Zhu L, Qian C, Tian H, Zhao Z, Jin L, Yang D. Antibacterial Activity of the Essential Oil From Litsea cubeba Against Cutibacterium acnes and the Investigations of Its Potential Mechanism by Gas Chromatography-Mass Spectrometry Metabolomics. *Front Microbiol*. 2022 Mar 2;13:823845. doi: 10.3389/fmicb.2022.823845. PMID: 35308342; PMCID: PMC8924494.
- [3] Wu N, Xing M, Chen Y, Zhang C, Li Y, Song P, Xu Q, Liu H, Huang H. Improving the productivity of malic acid by alleviating oxidative stress during *Aspergillus niger* fermentation. *Biotechnol Biofuels Bioprod*. 2022 Dec 29;15(1):151. doi: 10.1186/s13068-022-02250-7. PMID: 36581946; PMCID: PMC9801644.

References:

- [1] Esmail S, Kakey, Amez A, Ismael. Evaluation of Oxidative Stress Status in Aged Human in relation to some Diseases. *International Conference on Pure and Applied Sciences*. August 2018;

Related Products:

BC0920/BC0925 Fructose 1,6-bisphosphatase(FBP) Activity Assay Kit

BC3320/BC3325
BC3310/BC3315

Glucose-6-phosphatase Activity Assay Kit
Phosphoenolpyruvate Carboxykinase(PEPCK) Activity Assay Kit

