

Soil carbonic anhydrase (S-CA) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Cat No: BC5500

Size: 50T/24S

Components:

Reagent I: Liquid 75 mL×1, Storage at 2-8°C;

Reagent II: Powder×2, Storage at 2-8°C. Before use, take a bottle of Reagent II and add 1mL acetone to dissolve the powder thoroughly, then add 10mL distilled water. Unused reagents can be stored in aliquots at -20°C for 1 weeks, avoiding repeated freezing and thawing;

Standard solution: Liquid 1 mL×1, Storage at 2-8°C; 5μmol/mL phenol standard solution. Before use, 50μL of 5μmol/mL phenol standard solution was taken into the reagent bottle, and 750μL distilled water was added to mix thoroughly to form 0.3125μmol /mL phenol standard solution.

Product Description:

Carbonic Anhydrase (CA, EC4.2.1.1) is a metal enzyme with Zn²⁺ as the active center, which can be used to efficiently catalyze reversible hydration reaction of CO₂: $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{HCO}_3^- + \text{H}^+$, with a catalytic rate up to 10⁷ times that of natural conditions, which is one of the fastest enzymes known so far.

Carbonic anhydrase can catalyze the reaction of acetic acid to nitrophenyl ester to p-nitrophenol. The activity of carbonic anhydrase can be reflected by detecting the increase rate of absorption value at 405nm.

Required but Not Provided:

Ultraviolet spectrophotometer, desk centrifuge, water-bath/ constant temperature incubator, adjustable pipette, 30-50 mesh sieve, 1 mL quartz cuvette, Toluene, ice and distilled water.

Protocol

I. Preparation:

Fresh soil sample natural air dry or 37°C oven air dry, 30~50 mesh sieve.

II. Determination procedure:

1. Preheat ultraviolet spectrophotometer for 30 minutes, adjust wavelength to 405 nm, set the counter to zero with distilled water.
2. Preheat Reagent I in 37°C (mammal) or 25°C (other species) for 15 minutes.
3. Standard tube measurement
 - 1) Standard tube measurement: 320μL standard solution and 1180μL Reagent I were added into 2mL EP tube, thoroughly mixed, and the light absorption value at 405nm was measured in 1mL glass cupola, which was referred to as A_s.
 - 2) Standard blank tube measurement: 320μL distilled water and 1180μL Reagent I were added

into 2mL

EP tube. After full mixing, the absorbance value at 405nm was measured in 1mL glass cupola, which was recorded as A_{SB} .

3) calculation $\Delta A_S = A_S - A_{SB}$. (Standard tubes and standard blank tubes only need to be done 1-2 times.)

4. Operation table:

Reagent Name (μL)	Test tube (A_T)	Control tube (A_C)
Sample	0.2g	0.2g
Toluene	50	50
Fully shake the soil to make it moist, and leave it at room temperature for 15minutes		
Reagent I	1130	1130
		Boil for 10min and cool with ice water
Reagent II	320	320
After reaction at 37°C for 5minutes, it was immediately placed in an ice bath, and then centrifuged at 4°C for 15000g for 10minutes. The supernatant was absorbed into 1mL glass cupola to determine the light absorption value at 405nm, which was denoted as A_T and A_C . Calculate $\Delta A = A_T - A_C$. (Each measuring tube corresponds to a pair of tubes).		

III. Calculation of soil CA activity:

Sample weight:

Unit definition: The catalytic production of 1 μmol p-nitrophenol per g of soil per minute at 37°C was defined as a unit of enzyme activity.

$$S\text{-CA (U/g weight)} = C_S \times \Delta A \div \Delta A_S \times V_S \div W \div T \times F = 0.02 \times \Delta A \div \Delta A_S \div W \times F$$

C_S : Standard concentration, 0.3125 $\mu\text{mol/mL}$;

V_S : Standard solution volume added to the reaction system, 0.32mL;

T : Reaction time, 5 minutes;

W : Sample weight, g;

F : Sample dilution ratio.

Note:

- If $A > 1.5$ or $\Delta A > 0.8$, the sample size can be reduced or the enzymatic reaction time at 37°C can be shortened; When $\Delta A < 0.02$, the sample size can be increased or the enzymatic reaction time at 37°C can be prolonged. Note that the calculation formula is changed simultaneously during calculation.
- If the supernatant to be measured is still cloudy after centrifugation, try to increase the centrifugal speed or extend the time, for example, 20000g, centrifugation at 4°C for 10minutes.

Experimental example:

1. Take 0.2014g soil sample No. 65, operate according to the measurement steps, use 1mL glass cuvette to measure $\Delta A = A_T - A_C = 0.595 - 0.400 = 0.195$, $\Delta A_S = A_S - A_{SB} = 0.448 - 0 = 0.448$, put into the formula to calculate:
S-CA activity (U/g weight) = $0.02 \times \Delta A \div \Delta A_S \div W \times F = 0.043$ U/g weight

2. Take 0.2015g soil sample No. 1, operate according to the measurement steps, use 1mL glass cuvette to measure $\Delta A = A_T - A_C = 0.275 - 0.137 = 0.138$, $\Delta A_S = A_S - A_{SB} = 0.448 - 0 = 0.448$, put into the formula to calculate:
S-CA activity (U/g weight) = $0.02 \times \Delta A \div \Delta A_S \div W \times F = 0.031$ U/g weight

Recent Product Citations:

[1] Li W, Yu L J, Yuan D X, et al. A study of the activity and ecological significance of carbonic anhydrase from soil and its microbes from different karst ecosystems of Southwest China[J]. Plant and Soil, 2005, 272(1):133-141.

Related Products:

BC0150/BC0155	Soil Cellulase (S-CL) Activity Assay Kit
BC0160/BC0165	Soil β -glucosidase (S- β -GC) Activity Assay Kit

