

# Nitric Oxide Synthase Subtypes (TNOS, iNOS, cNOS) Activity Assay Kit

**Note:** It is necessary to predict 2-3 large difference samples before the formal determination.

**Operation Equipment:** Spectrophotometer/Microplate reader

**Cat No:** BC5695

**Size:**100T/48S

## Components:

**Extract solution I:** Liquid 60mL×1. Store at -20°C.

**Extract solution II:** Liquid 0.6mL×2. Store at -20°C. It is volatile and sealed immediately after use.

**Buffer solution:** Liquid 20mL×1. Store at 2-8°C.

**Reagent I:** Liquid 2.8mL×1. Store at 2-8°C.

**Reagent II:** Powder×1. Store at -20°C. It is in the glass bottle. Add 6mL Buffer solution into Reagent II. It can be stored at -20°C for 4 weeks after dispensing to avoid repeated freezing and thawing.

**Reagent III:** Liquid 30μL×1. Store at 2-8°C.

**Reagent III working solution:** Reagent III and Buffer solution are mixed by the ratio of 2μL: 198μL (0.2mL, 10T) to make Reagent III working solution according to sample number before use.

**Reagent IV:** Powder×1. Store at -20°C. Add 0.6mL Buffer solution into Reagent IV before use. It can be stored at -20°C for 4 weeks after dispensing to avoid repeated freezing and thawing.

**Reagent V:** Powder×1. Store at -20°C. It is in the glass bottle. Add 1.2mL Buffer solution into Reagent V. It can be stored at -20°C for 4 weeks after dispensing to avoid repeated freezing and thawing.

**Working solution:** Reagent I, Reagent II, Reagent III working solution and Reagent IV are mixed by the ratio of 0.2mL: 0.5mL: 0.2mL: 0.05mL (0.95mL, 10T) to make Working solution according to sample number before use.

**Reagent VI:** Liquid 0.6mL×1. Store at 2-8°C.

**Reagent VII:** Liquid 1.5mL×1. Store at 2-8°C.

**Reagent VIII:** Liquid 30μL×1. Store at 2-8°C.

**Reagent VIII working solution:** Reagent VIII and Buffer solution are mixed by the ratio of 5μL: 225μL (0.23mL, 23T) to make Reagent VIII working solution according to sample number before use.

**Chromogenic solution A:** Liquid 7mL×1. Store at 2-8°C.

**Chromogenic solution B:** Liquid 7mL×1. Store at 2-8°C.

**Chromogenic solution:** Chromogenic solution A and Chromogenic solution B are mixed by the ratio of 1: 1 to make Chromogenic solution according to sample number before use.

**Standard:** Liquid 1mL×1, 10μmol/mL sodium nitrite solution. Store at 2-8°C. Mix 10μL 10μmol/mL sodium nitrite solution and 990μL distilled water to prepare a 0.1μmol/mL standard solution before use.

**Note:** Reagent II, Reagent IV and Reagent V are lyophilized powder. The differences in the amount of

these powders may seem to be large, but the actual qualities are the same. It does not affect the detection.

## Product Description:

Nitric oxide synthase (NOS, EC 1.14.13.39, written as total NOS (TNOS) after) is a type of enzyme that catalyzes the synthesis of NO from L-arginine in living organisms. It mainly exists in various cells such as vascular smooth muscle, macrophages, endothelial cells, neurons, liver cells, and glomerular mesangial cells. According to the different dependence of its enzyme activity on calcium ions, it can be divided into structural NOS (constructive NOS, cNOS) and injury induced NOS (inducible NOS, iNOS). The former requires a certain concentration of calcium ions to activate, while the latter does not rely on exogenous calcium ions.

NOS catalyzes L-arginine, molecular oxygen and NADPH to form NO and NADP<sup>+</sup>. NO is easily oxidized to form NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup> in aqueous solution. Under acidic conditions, NO<sub>2</sub><sup>-</sup> and diazonium sulfonamide produce diazo compounds. The compounds could further couple with naphthyl vinyl diamine. The product has a characteristic absorption peak at 550 nm and its absorbance value can be measured to calculate NOS activity.

### Reagents and Equipment Required but Not Provided:

Spectrophotometer/Microplate reader, centrifuge, balance, transferpettor, mortar/homogenizer/cell ultrasonic crusher, micro glass cuvette/96 well plate, ice and distilled water.

### Procedure:

#### I. Sample preparation

- Tissue:** It is suggested that 0.2 g of tissue with 0.98 mL of Extract solution I and 0.02 mL of Extract solution II and fully homogenized on ice bath. Centrifuge at 12000g for 15 minutes at 4°C to remove insoluble materials and take the supernatant on ice for testing.
- Bacteria/Cells:** Collect bacteria/cells into the centrifuge tube, after centrifugation discard supernatant. It is suggested that add 0.98 mL of Extract solution I and 0.02 mL of Extract solution II to 10 million of bacteria/ cells. Use ultrasonication to split bacteria/cells (place on ice, ultrasonic power 200W, working time 3 seconds, interval 7 seconds, repeat for 5 minutes). Centrifuge at 12000g for 15 minutes at 4°C to remove insoluble materials and take the supernatant on ice for testing.
- Liquid samples:** Detect directly. Centrifuge before detecting if there are precipitation in the samples.

**Note:** Extract solution I and Extract solution II are mixed by the ratio of 0.98mL: 0.02mL to prepare according to sample number before use.

#### II. Determination

- Preheat spectrophotometer for 30 min, adjust the wavelength to 550 nm and set counter to zero with distilled water.
- Add reagents in 1.5ml EP tube as the following:

Reagent (μL)	Total NOS test tube(A <sub>T1</sub> )	iNOS test tube(A <sub>T2</sub> )	Standard tube(A <sub>S</sub> )	Blank tube(A <sub>B</sub> )
Sample	60	60	-	-
Working solution	95	95	-	-
Reagent V	20	-	-	-
Reagent VI	10	-	-	-
Buffer solution	-	30	-	-
Mix and react at 37°C for 60min, bath at 100°C for 5min and cool to room temperature. Centrifuge at 11000g for 10 minutes at 4°C and take all supernatant.			-	-
Supernatant	all supernatant	all supernatant	-	-
Reagent VII	10	10	-	-
Reagent VIII working solution	10	10	-	-
Mix and react at 37°C for 30min.			-	-
Standard solution	-	-	60	-
Distilled water	-	-	145	205
Chromogenic solution	100	100	100	100
Mix and react for 10min at room temperature. Take 200μL reaction solution in a 96 well plate. Detect the absorbance value at 550 nm and record as A <sub>T1</sub> , A <sub>T2</sub> , A <sub>S</sub> and A <sub>B</sub> . ΔA <sub>T1</sub> =A <sub>T1</sub> -A <sub>B</sub> . ΔA <sub>T2</sub> =A <sub>T2</sub> -A <sub>B</sub> . ΔA <sub>S</sub> =A <sub>S</sub> -A <sub>B</sub> . Blank tube and standard tube need to test once or twice.				

### III. NOS activity calculation:

#### 1. Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes production of 1 nmol NO in the reaction system per minute every milligram protein.

$$(1) \text{ TNOS activity (U/mg prot)} = (\Delta A_{T1} \div \Delta A_S \times C_S) \times V_S \div (V_S \times C_{Pr}) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T1} \div \Delta A_S \div C_{Pr} \times F$$

$$(2) \text{ iNOS activity (U/mg prot)} = (\Delta A_{T2} \div \Delta A_S \times C_S) \times V_S \div (V_S \times C_{Pr}) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T2} \div \Delta A_S \div C_{Pr} \times F$$

$$(3) \text{ cNOS activity (U/mg prot)} = \text{TNOS activity} - \text{iNOS activity}$$

#### 2. Sample mass:

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes production of 1 nmol NO in the reaction system per minute every gram sample.

$$(1) \text{ TNOS activity (U/g mass)} = (\Delta A_{T1} \div \Delta A_S \times C_S) \times V_S \div (W \times V_S \div V_E) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T1} \div \Delta A_S \div W \times F$$

$$(2) \text{ iNOS activity (U/g mass)} = (\Delta A_{T2} \div \Delta A_S \times C_S) \times V_S \div (W \times V_S \div V_E) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T2} \div \Delta A_S \div W \times F$$

$$(3) \text{ cNOS activity (U/g mass)} = \text{TNOS activity} - \text{iNOS activity}$$

#### 3. Bacteria/Cells number:

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes production of 1 nmol NO in the reaction system per minute every  $10^6$  bacteria/cells.

$$(1) \text{ TNOS activity (U/}10^6 \text{ cell)} = (\Delta A_{T1} \div \Delta A_S \times C_S) \times V_S \div (N \times V_S \div V_E) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T1} \div \Delta A_S \div N \times F$$

$$(2) \text{ iNOS activity (U/}10^6 \text{ cell)} = (\Delta A_{T2} \div \Delta A_S \times C_S) \times V_S \div (N \times V_S \div V_E) \times 10^3 \div T \times F = 1.67 \times \Delta A_{T2} \div \Delta A_S \div N \times F$$

$$(3) \text{ cNOS activity (U/}10^6 \text{ cell)} = \text{TNOS activity} - \text{iNOS activity}$$

#### 4. Liquid volume:

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes production of 1 nmol NO in the reaction system per minute every milliliter liquid sample.

$$(1) \text{ TNOS activity (U/mL)} = (\Delta A_{T1} \div \Delta A_S \times C_S) \times V_S \div V_S \times 10^3 \div T \times F = 1.67 \times \Delta A_{T1} \div \Delta A_S \times F$$

$$(2) \text{ iNOS activity (U/ mL)} = (\Delta A_{T2} \div \Delta A_S \times C_S) \times V_S \div V_S \times 10^3 \div T \times F = 1.67 \times \Delta A_{T2} \div \Delta A_S \times F$$

$$(3) \text{ cNOS activity (U/ mL)} = \text{TNOS activity} - \text{iNOS activity}$$

$C_S$ : sodium nitrite concentration of standard solution,  $0.1 \mu\text{mol/mL}$ ;

$V_S$ : Add sample supernatant volume, 0.06 mL;

$V_E$ : Add volume of Extract solution I and Extract solution II, 1mL;

$C_{pr}$ : Sample protein concentration, mg/mL;

$W$ : Sample mass, g;

$N$ : Cell amount,  $10^6$  for one unit;

$10^3$ : Unit conversion factor,  $1 \mu\text{mol} = 10^3 \text{ nmol}$ ;

$T$ : Reaction time, 60 min;

$F$ : Dilution factor

#### Note:

1. NOS is not stable and easy to inactivate. It is recommended to use fresh samples for experiments and stored at  $-20^\circ\text{C}$  for fresh samples if not tested immediately.
2. The prepared Reagent II need to be stored at  $-20^\circ\text{C}$  after taking out the required volume according to the sample number.
3. If  $\Delta A_T$  is less than 0.005, or  $A_T$  is closed to  $A_B$ , it is recommended to increase added sample supernatant volume or prolong time of the first reaction at  $37^\circ\text{C}$  before determination. If  $\Delta A_T$  is more than 0.5, it is recommended to dilute the sample with Buffer solution before determination. And modify the calculation formula.

#### Experimental example:

1. Take 0.2047g fresh mouse liver samples, add 0.98 mL of Extract solution I and 0.02 mL of Extract solution II, grind the homogenate with ice bath. Then operate according to the determination steps, using 1mL glass cuvette and calculate  $\Delta A_{T1} = A_{T1} - A_B = 0.114 - 0.045 = 0.069$ ,  $\Delta A_{T2} = A_{T2} - A_B = 0.072 - 0.045 = 0.027$ ,  $\Delta A_S = A_S - A_B = 0.550 - 0.045 = 0.505$ . The result is calculated according to the



sample mass:

$$\text{TNOS activity (U/g mass)} = 1.67 \times \Delta A_{T1} \div \Delta A_S \div W \times F = 1.115 \text{ U/g}$$

$$\text{iNOS activity (U/g mass)} = 1.67 \times \Delta A_{T2} \div \Delta A_S \div W \times F = 0.436 \text{ U/g}$$

$$\text{cNOS activity (U/g mass)} = \text{TNOS activity} - \text{iNOS activity} = 0.679 \text{ U/g}$$

2. Take 60 $\mu$ L horse serum and operate according to the determination steps, using 1mL glass cuvette and calculate  $\Delta A_{T1} = A_{T1} - A_B = 0.066 - 0.045 = 0.021$ ,  $\Delta A_{T2} = A_{T2} - A_B = 0.062 - 0.045 = 0.017$ ,  $\Delta A_S = A_S - A_B = 0.550 - 0.045 = 0.505$ . The result is calculated according to the sample mass:

$$\text{TNOS activity (U/mL)} = 1.67 \times \Delta A_{T1} \div \Delta A_S \div W \times F = 0.069 \text{ U/mL}$$

$$\text{iNOS activity (U/mL)} = 1.67 \times \Delta A_{T2} \div \Delta A_S \div W \times F = 0.056 \text{ U/mL}$$

$$\text{cNOS activity (U/mL)} = \text{TNOS activity} - \text{iNOS activity} = 0.013 \text{ U/mL}$$

### References:

[1] List BM, Klösch B, Völker C. et al. Characterization of bovine endothelial nitric oxide synthase as a homodimer with down-regulated uncoupled NADPH oxidase activity: tetrahydrobiopterin binding kinetics and role of haem in dimerization[J]. Biochemical Journal, 1997, 323(1): 159-165.

[2] Dawson J, Knowles RG. A microtiter-plate assay of nitric oxide synthase activity[J]. Molecular Biotechnology, 1999, 12(3): 275-279.

[3] Förstermann U, Sessa WC. Nitric oxide synthases: regulation and function[J]. European Heart Journal, 2012, 33(7): 829-837.

[4] Kazakov A, Hall R, Jagoda P. et al. Inhibition of endothelial nitric oxide synthase induces and enhances myocardial fibrosis [J]. Cardiovascular Research, 2013, 100(2):211-221.

### Related Products:

BC1470/BC1475	Nitric Oxide (NO) Content Assay Kit
BC5480/BC5485	Nitric Oxide (NO) Content Assay Kit
BC5680/BC5685	Nitric Oxide Synthase (NOS) Activity Assay Kit